

Bacillus intermedius ribonuclease as inhibitor of cell proliferation and membrane current

Olga Ilinskaya ^a, Katrin Decker ^b, Andreas Koschinski ^b, Florian Dreyer ^b,
Holger Repp ^{b,*}

^a Department of Microbiology, Kazan State University, Kremlevskaya Street 18, 420008 Kazan, Russia

^b Rudolf-Buchheim-Institute of Pharmacology, Justus-Liebig-University of Giessen, Frankfurter Street 107, 35392 Giessen, Germany

Received 3 May 2000; received in revised form 6 June 2000; accepted 6 September 2000

Abstract

The antiproliferative action of the guanine-specific ribonuclease secreted by *Bacillus intermedius* (binase) was studied in different chicken and mouse cell lines. The proliferation rate of chicken embryo fibroblasts, either normal or Rous sarcoma virus-transformed, was significantly reduced by binase treatment. Among mouse fibroblasts, *v-ras*-transformed NIH3T3 cells were sensitive to binase, whereas the growth of non-transformed, *v-src*-transformed or *v-fms*-transformed NIH3T3 cells was not affected. A 48 h treatment with binase inhibited the Ca^{2+} -dependent K^{+} current of *v-ras*-transformed NIH3T3 cells but had no effect on this membrane current in non-transformed and in *v-src*- or *v-fms*-transformed NIH3T3 cells. Our results suggest that mammalian cells expressing the *ras*-oncogene are a potential target for the antiproliferative action of binase. © 2001 Elsevier Science Ireland Ltd. All rights reserved.

Keywords: *Bacillus intermedius* RNase (binase); Chicken embryo fibroblasts; Mouse fibroblasts; Transformation; Proliferation; Ca^{2+} -dependent K^{+} current

1. Introduction

A number of ribonucleases, such as binase (*Bacillus intermedius* RNase), pancreatic RNase

A, bovine seminal ribonuclease, and RNase from oocytes of *Rana pipiens* (onconase) were shown to possess enzymatic and biological activities which may be exploited for the development of novel cytotoxic agents (Youle et al., 1993; Di Donato et al., 1994; Mikulski et al., 1995; Boque and Wlodawer, 1996; Prior et al., 1996). To assess the possibility of a future clinical application of bacterial ribonucleases, the toxicity of these enzymes in various cell lines is currently under investigation. Compared to pancreatic RNase A, binase is a more effective growth inhibitor of Ehrlich ascites tumor cells and also of the ascitic form of lymph-

Abbreviations: CEF, chicken embryo fibroblast; EGTA, ethyleneglycol-bis(β -aminoethyl)-*N,N,N',N'*-tetraacetic acid; HEPES, 2-[4-(2-hydroxyethyl)-1-piperazine]-ethanesulfonic acid; LPA, lysophosphatidic acid; RNA, ribonucleic acid; RNase, ribonuclease; RSV, Rous sarcoma virus.

* Corresponding author. Tel.: +49-641-9947622; fax: +49-641-9947609.

E-mail address: holger.repp@pharma.med.uni-giessen.de (H. Repp).